

Periglacial phenomena in the high mountains of northwestern Argentina

Ana Lia Ahumada*

Periglacial processes and features occur at high elevations in the mountains of the Sierras Subandinas, Sierras Pampeanas, Cordillera Oriental, Puna and Cordillera del Límite of Argentina (22–28°S, 65–68°W). This paper outlines our knowledge of these periglacial processes and landforms, as reported in the literature over the last 80 years. Landforms investigated include active and inactive rock glaciers, patterned ground, solifluction, proglacial lakes and other processes such as mineral segregation and concentration. Fossil periglacial landforms are an indication of past climates. Knowledge of geocryological phenomena in the northwest of Argentina has contributed to a better understanding of climate change.

Background

The northwest of Argentina is an almost unexplored region that has been little studied in terms of its periglacial and geocryological characteristics. However, its geographic position and its significance for human settlement during the Quaternary call for a detailed analysis of its geomorphological and palaeoclimatic record.

The region is located between 22° and 28°S and from c. 65° to 68°W, covering four distinctive, morphostructurally different regions, which have a N–S orientation and increase in altitude towards the west. From west to east these are: the Puna de Atacama, the Sierras Subandinas, the Cordillera Oriental, and the Pampean Sierras of the north (Fig. 1).

The high mountains and relief areas are of particular importance for their effect on climate and the drainage system. The Cordillera del Límite, at the western limit of this region, is characterized by the presence of volcanic craters of Tertiary and Quaternary age, some of them now extinct. The highest summits range between 5200 m and almost 6900 m a.s.l. (Fig. 1). The most important summits are the Cerro Panizos (5259 m a.s.l.), the Cerro Vilama (5678 m), the Nevado San Pedro (5750 m), the Volcán Socompa (6301 m), the Cerro Llullaillaco (6723 m), the Volcán del Azufre (5680 m), the Cerro del Laudo (6400 m), the Cerro de Incahuasi (6620 m) and the Volcán Ojos del Salado (6885 m). The eastern limit, also called Prepuna, comprises the Cordillera Oriental, with the summits ranging between 4200 m and almost 6400 m: the Sierras Santa Victoria (Cerro Negro, 5029 m) and Zenta, which continue towards the south as the Sierras de Chañi (Nevado de Chañi, 6200 m), Acay (5950 m), Nevados del Palermo (6120 m), Nevados de Cachi (6380 m), Nevados de Catreal and Nevados de Chuscha (5468 m), the extensions of which combine with the extensions of the Sierras Pampeanas in the north; Cumbres Calchaquies, Sierra del Cajón or Sierra de Quilmes (Cerro Negroara, 4200 m) and further south, the Sierra del Aconquija with the Cordón de las Animas and the Nevados del Aconquija (Morro del Zarzo,

5064 m; Cerro Negro, 4700 m; Cerro del Bolsón, 5500 m and Nevado del Candado, 5450 m). These mountain ranges or belts, with an average height of over 5000 m, converge in the Puna region, a high-altitude plateau with an average elevation of over 3500 m, with salt pans, an arid climate and a markedly reduced humidity. The orographic belts lie above the 0°C isotherm (4600 m a.s.l.) but, at present, hardly any evidence of glaciers is found. During the Pleistocene, however, these belts were covered by glaciers at least three times.¹

In this geographical setting the climatic changes during the Quaternary, and in particular during the Late Pleistocene and Holocene, left geomorphological evidence that may be associated with the intense action of a periglacial environment, taking into account the region's altitude with pronounced annual temperature variations and strong mechanical weathering, only one of many indicators of a periglacial high mountain environment.

Numerous authors since the beginning of the last century have referred to present and past periglacial conditions in the northwest of Argentina. Their observations are summarized here.

Climate setting

Owing to its proximity to the Tropic of Capricorn, the study area experiences a warm climate with little seasonal variation of daylight hours. However, the altitudinal diversity, the N–S orientation of the mountain ranges as well as its continentality are responsible for a complex mosaic of climatically different regions, temperature variations depending on altitude, and a variable distribution of mean annual precipitation, with a pronounced rainfall peak in summer. This creates extreme situations: very warm and humid climates in the eastern valleys and cold deserts on the plateau of the Puna region.

Rainfall is intense in summer, with a marked east to west gradient, and is associated with easterly winds from the Atlantic which penetrate as NE flows into the mountain reliefs. On the eastern borders of the mountains, orographic precipitation falls at two main levels: the one at an altitude of 1500–1800 m with annual precipitation between 1000 and 2000 mm, with a steep decline towards the west until it reaches values of less than 100 mm in the Puna region (some areas receive at little as 50 mm).² In some cases, in the Sierra del Aconquija and at Cumbres Calchaquies, the first orographic barrier to the easterly winds, mean annual precipitation of 2500 mm (at 2500 m) has been recorded.³ Above this level, rainfall declines on the eastern slopes of these mountains to 600–500 mm/yr.²

In the Puna region the mean annual air temperature (MAAT) is less than 5°C, with a large daily amplitude. The mean annual maximum and minimum temperatures are 16°C and –4°C, respectively. An occasional very strong effect is caused by the so-called 'white winds'. They cause snowstorms produced by the entry of westerly winds from the Pacific, which at heights above 5000 m penetrate the area in winter. In the Puna region, winds blow predominantly from the west; their frequency and intensity increase slightly during the dry winter season. Humid easterly winds blow in summer. Despite their relatively rare

*CONICET–Fundación Miguel Lillo, Miguel Lillo 251, 4000- San Miguel de Tucumán, Argentina. E-mail: iesglount@infovia.com.ar

occurrence, the easterly winds bring 88–96% of the area's precipitation.⁴ In summary, the Puna region receives precipitation mainly during the summer, with a general decrease towards the west. The evapotranspiration is high, with the result that the mean annual relative humidity is less than 40%. The global annual radiation is over 200 kcal/cm², according to Minetti.²

Meteorological data for this part of the country are sparse and concentrated on the eastern slopes of the valleys. Weather stations in the Puna region are scarce and the few existing ones have records of a few years only. To obtain a better idea of the climatic conditions which favour a periglacial environment in a high mountain area, more climatic data will have to be collected, so that the circumstances that enhance its presence can be defined more precisely. The available climatic data for the region have been compiled by Minetti.²

Periglacial regions in the northwest of Argentina

The relation between mountain permafrost and climate is poorly understood. Mountain permafrost is the result of a complex interaction of environmental factors, the most important of which is climate. The general climatic conditions in high mountain areas depend mainly on latitude, height and continentality and, less importantly, on local conditions which produce variations on a limited geographical scale.

In high mountain areas the periglacial environment may be classified into different zones as a function of altitude. Gorbunov⁵ introduced a classification of relations between permafrost/soil and temperature/vegetation based on an index of continentality. The latter is based on the difference between the elevation of the glaciers' equilibrium line and the lowest permafrost limit. This index allows us to distinguish five different types of permafrost along the Andes: the Equadorian type, the Himalayan type, the Central Andean type, the Tibetan type, and the New Zealand type. The study area belongs to the Central Andean type.

Corte,⁶ in his map of current geocryological processes in Argentina, established the lower permafrost limit at 4000 m at the latitude of Salta, following a method similar to Gorbunov's, as part of a periglacial inventory of the area.

Garleff and Stingl⁷ argued for a 9°C depression of the mean annual air temperature during the Pleistocene for the Puna region. On the evidence of geomorphological data, among others, the corresponding descent of so-called 'almost continuous permafrost' and strong cryogenic activity is estimated to have been 1000 m in the NW of Argentina.^{8,9}

Glacial and periglacial processes and phenomena

Table 1 summarizes the glacial and periglacial processes and landforms in the northwest of Argentina that have been

Table 1. The glacial and periglacial processes and landforms in the northwest of Argentina reported in the literature.

Site (range) Mountain name Peak name	Reference	Summit elevation (m a.s.l.)	Periglacial processes and landforms
Puna	14	4000	Rock glaciers
Puna	15, 16	4000	Rock glaciers
Nevado de Queva	18	6130	Rock glaciers
Co Granadas	18	5705	Rock glaciers
Acay	20	4500	Rock glaciers
Nevados de Cachi	20	4500	Rock glaciers
Sierra de Santa Victoria	21, 22	4500	Rock glaciers
Sierra del Aconquija	23, 24, 26, 28	4000	Rock glaciers
Nevados de Chuscha	11	4200–4700	Rock glaciers
Mina Pirquitas	21	4000	Frost depth
Vn Ojos del Salado	27	6885	Frost depth
Puna	9	4000	Rock weathering
Puna	14	4000	Rock weathering
Puna	15, 16	4000	Rock weathering
Puna	18	4000	Rock weathering
Vn Tuzgle	29	5480	Sorted polygons.
Vn Socompa	29	6031	Sorted polygons.
Cumbres Calchaquies	4	4200	Sorted polygons.
27°S	30	2600	Solifluction
Valle de Tafi	31	–	Solifluction
El Rincón	26, 28, 32	2500	Solifluction
Puna	14	–	Cryoplains
Cumbres Calchaquies	33	4200–4700	Cryoplains
Cumbres Calchaquies	4	4200	Needle ice
Nacientes del Río Los Reales	28	2800	Needle ice
Sierras del Aconquija Cumbres Calchaquies	34	–	Mass wasting
Sierras del Aconquija Cumbres Calchaquies	31	–	Mass wasting
Sierras del Aconquija Cumbres Calchaquies	12	–	Mass wasting
Sierras del Aconquija Cumbres Calchaquies	33	Over 4000	Peatlands
Nacientes del Río Los Reales	26	2800	Peatlands
Cumbres Calchaquies y Sierra del Aconquija	12	–	Proglacial lakes
Cordillera Oriental	35	–	Proglacial lakes

reported in the literature of the last 80 years.

Present-day glacial conditions. There are several records of glacial cirque floors; Penck¹⁰ determined contemporary glacial cirque floors at 5000 m. The average height of these features on the eastern slopes of the Sierra Aconquija is 4300 m.¹ On the western slopes of the Aconquija, glacial cirque floors are formed at over 4400–4600 m.¹¹ In the Sierra de Quilmes and Nevados de Chuscha (5468 m), glacial cirques are at 4200–4700 m and are now occupied by several small rock glaciers.¹² In Cumbres Calchaquies, there are glacial features at Quebrada del Matadero (c. 4700 m) and in the Alto de la Mina Mt (c. 4700 m), in the Laguna de Huaca Huasi area.^{13,14}

Rock glaciers. Keidel's studies on Puna¹⁵ were the first to describe block or rock slopes as characteristic of the region. He probably included rock glaciers in these deposits. He described their lower levels up to 4000 m a.s.l., did not observe any activity, but recognizes their palaeoclimatic importance and considers them an example of morphology characteristic of cold deserts.

Catalano,^{16,17} in ref. 6, referred to glacial action in Puna and pointed out the presence of rocks agglomerated by ice and flowing under gravity as a glacier. He named these forms litho glaciers.

Igarzábal¹⁸ noticed the presence of an ice nucleus covered by detritus in the Nevado del Queva (6130 m) and Granadas Mt (5707 m), that fed permanent streams. He called them glaciolithic deposits, using Corte's classification of rock glaciers.¹⁹

Igarzábal²⁰ compiled an inventory of active and inactive rock glaciers in the Acay region and the Nevados de Cachi, and determined the lower limit of rock glaciers at 4500 m.

In the Sierra de Santa Victoria, Corte *et al.*²¹ observed active rock glaciers at 4500 m a.s.l. In the same region Zipprich *et al.*²² distinguished three generations of rock glaciers, with presently active glaciers, and determined the lower limit of periglacial

processes at 700 m, at a temperature of 6°C, corresponding to c. 27 980 ± 190 BP.

Ahumada²³ stressed the importance of setting high basins in the Sierras Pampeanas in order to detect rock glaciers that feed local rivers. She compiled an inventory of active and inactive rock glaciers of the northernmost region of Sierra del Aconquija.^{24, 25} Figure 2 shows the location of rock glaciers in Cordón de las Ánimas, on the eastern slopes of the Sierra del Aconquija.²⁶

Frost-depth measurements. Corte *et al.*²¹ measured the freezing depth in Pirquitas mine (4000 m a.s.l.) and found a stable temperature of 0°C at a depth of 1 m. This observation corresponds to a depth of 1.30 m, according to thermistor readings. Cobos and Corte²⁷ registered a 70-cm-deep seasonal frost in permafrost at 4700 m, on the Volcán Ojos del Salado.

Needle ice. Halloy⁴ described the action of needle ice on the grassland around Huaca Huasi lagoon, in Cumbres Calchaquíes, at 4000 m a.s.l. In a nearby region, at a lower elevation, Ahumada *et al.*²⁸ described the action of needle ice as leaving a rake-like pattern on the ground, on slopes of less than 5° at 2500 m, indicating diurnal freezing.

Cryoweathering. Penck,¹⁰ Keidel¹⁵ and Catalano¹⁶ attributed the mechanical weathering phenomenon in Puna to the great daily and seasonal temperature variation. Igarzábal¹⁸ reported the extensive cryoweathering of Puna's land surface, that causes generalized surface erosion.

Patterned ground. Corte²⁹ identified patterned ground associated with the Tuzgle (5480 m) and Socompa (6031 m) volcanoes. Halloy⁴ found small areas of patterned ground in Cumbres Calchaquíes, at 4000 m.

Solifluction. Auer³⁰ reported that, at 27°S, the solifluction limit has descended 2000 m and is now at 2600 m. Sayago and Collantes³¹ described solifluction processes in the smoothing of the Tafi Valley's slopes. Ahumada *et al.*^{26,28,32} described solifluction grounds at El Rincón, Sierra del Aconquija, in a seasonal freeze-thaw environment, between 2500 and 4000 m a.s.l.

Cryoplanation. Keidel¹⁵ was one of the first to mention cryoplains and he called them 'summit' cryoplanation surfaces. Though he did not indicate specific sites, he recognized their palaeoclimatic importance in Puna.

In the Sierras del Aconquija and Cumbres Calchaquíes, at over 4000 m, Halloy *et al.*³³ described a high plain modified by glacier and periglacial action, with beach-like terraces.

Mass wasting. The big clastic deposits surrounding the Sierras del Aconquija and Cumbres Calchaquíes have been described in the literature by Bossi³⁴ and Porto and Danieli³⁵ as fanglomeratic deposits accumulated by alluvial and/or periglacial activity.

Sayago and Collantes³¹ and Sayago *et al.*¹³ described the large cenoglomeratic clastic accumulations over the covered glacies of the eastern slopes of the Sierra del Aconquija and the Cumbres Calchaquíes. They concluded that they originated as deposits under gravity at the same time as the glacial episodes detected in the peak area, which they considered to be of periglacial origin.

Peatlands. Halloy *et al.*³³ noted the existence of mountain peatlands under conditions of seasonal freezing in the high peaks of the Sierra del Aconquija and Cumbres Calchaquíes.

Ahumada *et al.*²⁶ described peatland deposits at 2800 m, in the Reales River basin, in a seasonal freezing environment.

Proglacial lakes. Proglacial lake sequences have been described by Sayago and Collantes³¹ and Sayago *et al.*¹³ in the Mollar region, Sierra del Aconquija. Ahumada and Vides³⁶ deduced periglacial conditions in the lake deposits of Cordillera Oriental.

Chemical segregation in freezing conditions. Ahumada and Vides³⁶ identified mirabilite (Na₂SO₄·10H₂O) by X-ray diffraction in lake




Fig. 2. The location of rock glaciers in Cordón de las Ánimas.

sequences of the southern region of Cordillera Oriental. The presence of hyper-hydrated sulphate indicates a cooling of the water due to the proximity of glaciers. Mirabilite³⁷ is produced by the segregation of pure sodium sulphate crystals. These deposits are evidence for an active periglacial environment, and were dated by Trauth and Strecker³⁸ as 35 650 ± 380 yr BP.

Heavy minerals concentration. Ahumada and Heinrich³⁹ analysed the Pb, Sn and Ag content of 230 drillings in the alluvium terraces at Pirquitas mine in Puna. The metals were concentrated mainly at a depth of 1.10–1.30 m. They attributed these concentrations to the seasonal freezing and thawing that took place after mineral deposition during the Late Pleistocene.

Altitudinal zonation. Garleff and Stingl⁷ defined the regional characterization of periglacial activity in relation to altitude. More recently, Ahumada *et al.*²⁶ established the elevation of periglacial processes in the Reales River basin (Sierra del Aconquija), and distinguished two levels: a lower level, from 2500 to 4000 m, with seasonal frosts, needle ice action and solifluction; and an upper level, from 4000 to an average of 4500–5000 m, with intense gelifraction, active and inactive rock glaciers, talus, old moraine deposits, and glacial cirques.

Conclusions

Many periglacial landforms have been reported from the high mountains of northwestern Argentina. They include rock glaciers, sorted polygons, frost weathering, solifluction, lakes, and periglacial mass wasting. Some periglacial phenomena, such as rock glaciers, solifluction, and frost weathering are active today. No investigations involving instrumented measurements of actual conditions have been conducted. Several authors deduced, however, that during the Last Glacial Maximum the altitude of periglacial activity descended between 1000 and 700 m below the present, corresponding to a temperature decrease of approximately 9°C to 6°C.

Among the questions to be addressed regarding periglacial phenomena in northwestern Argentina are the following:

- What is the extent and character of large-scale landforms in the high mountains?
- How much did climate permit the development of such geological features in a subtropical region?
- What is the relationship between the nature and altitude of fossil periglacial phenomena and past climate?

Improved knowledge of the character of the periglacial environment in this region will contribute to the development of models for sustainable development. These models may assist in the protection against natural disasters caused by anthropogenic interference in the environment.

This work was carried out with the support of the Miguel Lillo Foundation and of CONICET. I acknowledge the logistical assistance of geologists G. Ibáñez and M. Jiménez, and thank D. Tromboto for comments on the manuscript.

1. Fox A. and Strecker M. (1991). Pleistocene and modern snowlines in the Central Andes (24–28°S). In *Südamerika Geomorphologie und Paläoökologie im jüngeren Quartär*, eds K. Garleff and H. Stingl. *Bamberger Geographische Schriften* 11, 155–168.
2. Minetti J.L. (1999). *Atlas Climático del NW Argentino. Laboratorio Climatológico sudamericano*. Fundación Carl c. Zon Caldenius. Sede NOA.
3. Rohmeder W. (1943). Observaciones meteorológicas en la región encumbrada de las Sierras de Famatina y del Aconquija (República Argentina). *Anales de la Sociedad Científica Argentina* 136, 97–124.
4. Halloy S. (1985). *Climatología y edafología de alta montaña en relación con la composición y adaptación de las comunidades bióticas con especial referencia a las Cumbres Calchaquíes, Tucumán*. Ph.D. thesis. Univ. Microfilm International Pub. 85-02967. Ann Arbor, Michigan.
5. Gorbunov A. (1978). Permafrost investigations in high-mountain regions. *Arctic and Alpine Res.* 10, 283–294.
6. Corte A.E. (1982). Geocriología Argentina general y aplicada. *Revista del Instituto de Ciencias Geológicas* 5, 87–120.
7. Garleff K. and Stingl H. (1983). Hangformen und Hangformung in der periglazialen Höhenstufe der argentinischen Anden zwischen 27° und 55° südlicher Breite. In *Mesoformen des Reliefs im heutigen Periglazialraum*, eds H. Poser and E. Schunke, pp. 425–434. Göttingen.
8. Garleff K. and Stingl H. (1985). Höhenstufen und ihre raumzeitlichen Veränderungen in den argentinischen Anden. *Zbl. Geol. Paläont. Teil I*, H.11/12, 1701–1707.
9. Garleff K., Schäbitz F., Stingl H. and Veit H. (1991). Jungquartäre Landschaftsentwicklung und Klimageschichte beiderseits der Ariden Diagonale Südamerikas. In *Südamerika Geomorphologie und Paläoökologie im jüngeren Quartär*, eds K. Garleff and H. Stingl. *Bamberger Geographische Schriften* 11, 359–394.
10. Penck W. (1920). Der Südrand der Puna de Atacama (NW. Argentinien). Ein Beitrag zur Kenntnis des andinen Gebirgstypus und zur Frage der Gebirgsbildung. *Abh. Math. Phys. Kl. Sachsischen Akad. Wiss.* 37, 1–420.
11. Tapia A. (1925). Apuntes sobre el glaciario pleistocénico del Nevado del Aconquija. *Anales de la Sociedad Argentina de Geografía* 1, 313–365.
12. Strecker M.R. (1987). *Late Cenozoic landscape development, the Santa María valley, northwestern Argentina*. Ph.D. thesis, Cornell University, Ithaca.
13. Sayago J.M., Collantes M.M. and Arcuri C.B. (1991). El glaciario Finipleistoceno-Holoceno y su relación con los depósitos clásticos pedemontanos en la región montañosa de Tucumán. In *Südamerika Geomorphologie und Paläoökologie im jüngeren Quartär*, eds K. Garleff and H. Stingl. *Bamberger Geographische Schriften* 11, 155–168.
14. Arcuri C.B. (1988). Geomorfología y evolución paleoambiental de la zona de Huaca Huasi (Cumbres Calchaquíes, Tucumán, Argentina). *Seminario Facultad de Ciencias Naturales de la Universidad Nacional de Tucumán*.
15. Keidel J. (1922). Sobre la influencia de los cambios climáticos cuaternarios en el relieve de la región seca de los Andes centrales y septentrionales de la Argentina. *Boletín de la Dirección General de Minas, Geología e Hidrología, Publicación* 5, 3–19, Buenos Aires.
16. Catalano L.R. (1927). Datos hidrológicos del Desierto de Atacama. *Boletín de la Dirección General de Minas, Geología e Hidrología. Publicación* 35, 1–55.
17. Catalano L.R. (1930). Puna de Atacama (Territorio de Los Andes) Reseña Geológica y Geográfica. *Universidad Nacional del Litoral. Publicación* 8, 1–98.
18. Igarzábal A. (1982). El relieve de la Puna argentina. *Revista del Instituto de Ciencias Geológicas* 5, 45–66.
19. Corte A.E. (1976). Rock glaciers. *Biuletyn Periglacialny* 26, 157–197.
20. Igarzábal A. (1983). El sistema glaciolítico de la Cuenca Superior del Río Juramento, Provincia de Salta. *VIII Congreso Geológico Argentino, Actas* IV, 167–183.
21. Corte A.E., Tromboto D.T.A. and Ahumada A.L. (1982). *Relevamiento de la geomorfología criogénica del NW Argentino*. IANIGLA-CRICYT, Mendoza.
22. Zipprich M., Reizner B., Veit H., Zech W. and Stingl H. (1998). Upper Quaternary climate and landscape evolution in the Sierra de Santa Victoria (Cordillera Oriental, northwestern Argentina) deduced from geomorphologic and pedologic studies. *Terra Nostra* 5, 180–181.
23. Ahumada A.L. (1995). El periglacial de montaña en Sierras Pampeanas noroccidentales. In II seminario sobre Las Geociencias y el Cambio Global. *Asociación Geológica Argentina. Serie D: Publicaciones especiales* 2, 6.
24. Ahumada A.L. (1999). Los glaciares de escombros en las altas montañas del NW de Argentina. In *XIV Congreso Geológico Argentino. Resúmenes. Actas* I, 80. Salta.
25. Ahumada A.L. (2000). Periglacial phenomena in the NW of Argentina. In *31st International Geological Congress, Symposium on History of Geosciences*. CD edition.
26. Ahumada A.L., Ibáñez G. and Jiménez M.J. (in press). Pisos altitudinales de procesos geomórficos en la Cuenca del Río Los Reales, Monteros, Tucumán, Argentina. *Acta geológica lilloana*.
27. Cobos D. and Corte A. (1990). Geocryological observations in Ojos del Salado, Central Andes, Lat. 27°. IGCP/UNESCO Project 297, *Second Meeting, Geocryology of Southern Africa, Abstracts of papers*. Rhodes University, Grahamstown.
28. Ahumada A.L., Tromboto D.T.A., Ibáñez Palacios G. and Jiménez M.J. (2000). Guía de Campo. *Curso de Postgrado 'Los Fenómenos Periglaciales. Identificación, Determinación y Aplicación'*. Universidad Nacional de Tucumán.
29. Corte A.E. (1955). Contribución a la morfología periglacial especialmente criopedología de la República Argentina. *Acta fennica* 14, 83–102.
30. Auer V. (1970). The Pleistocene of Fuego-Patagonia; Par V, Quaternary problems of southern South America. *Anal. Acad. Sci. Fennicae* 100, 194.
31. Sayago J.M. and Collantes M.M. (1991). Evolución paleogeomorfológica del valle de Tafi (Tucumán, Argentina) durante el Cuaternario superior. In *Südamerika Geomorphologie und Paläoökologie im jüngeren Quartär*, eds K. Garleff and H. Stingl. *Bamberger Geographische Schriften* 11, 109–124.
32. Ahumada A.L., Ledesma F.C., Ibáñez Palacios G., Delgado M. and Jiménez M.J. (2000). Una evaluación de riesgo potencial en la región NE de la Sierra del Aconquija Tucumán, Argentina. *IX Congreso Geológico Chileno. Actas*, vol. 1: 16–20.
33. Halloy S., González J.A. and Grau A. (1994). Proyecto de creación del Parque Nacional Aconquija (Tucumán-Argentina). Informe N° 4. Fundación Miguel Lillo. *Serie Conservación de la Naturaleza* 9.
34. Bossi G.E. (1969). Geología y estratigrafía del sector S del Valle de Choromoro. *Acta geológica lilloana* 10, 17–64.
35. Porto J.C. and Danieli C.A. (1980). Hoja Trancas (provincias de Tucumán y Salta). Servicio Minero Nacional (unpubl. doc.).
36. Ahumada A.L. and Vides M.E. (2000). Mirabilia en secuencias lacustres del Pleistoceno tardío del NW argentino. *IX Congreso Geológico Chileno. Actas*, vol. 1: 586–589.
37. Grossman I.G. (1966). Origin of the sodium sulfate deposits of the Northern Great Plains of Canada and the United States. *U.S. Geological Survey Professional Paper* 600-B: B104–B109.
38. Trauth M.H. and Strecker M.R. (1999). Formation of landslide-dammed lakes during a wet period between 40,000 and 25,000 yr BP in northwestern Argentina. *Palaeogeog., Palaeoclimatol. Palaeocol.* 153, 277–287.
39. Ahumada A.L. and Heinrich S. (1986). Procesos de congelamiento y descongelamiento en la concentración Pb, Ag y Sn en el aluvión de Mina Pirquitas. IANIGLA-CRICYT, Mendoza (unpubl. doc.).